



DAMA/LIBRA results and perspectives

2011 MUNICH
TAUP


A.Incicchitti
INFN Roma

Roma2,Roma1,LNGS,IHEP/Beijing


- + by-products and small scale expts.: INR-Kiev
- + neutron meas.: ENEA-Frascati
- + in some studies on $\beta\beta$ decays (DST-MAE project): IIT Kharagpur, India



DAMA:an observatory for rare processes @LNGS



Relic DM particles from primordial Universe



(& invisible axions, ν 's)

&


Right halo model and parameters?



Some direct detection processes:

- Scatterings on nuclei

→ detection of nuclear recoil energy



- Inelastic Dark Matter: $W + N \rightarrow W^* + N$

→ W has Two mass states χ^+ , χ^- with δ mass splitting

→ Kinematical constraint for the inelastic scattering of χ^- on a nucleus


$$\frac{1}{2} \mu v^2 \geq \delta \Leftrightarrow v \geq v_{thr} = \sqrt{\frac{2\delta}{\mu}}$$

- Excitation of bound electrons in scatterings on nuclei

→ detection of recoil nuclei + e.m. radiation


- Conversion of particle into e.m. radiation

→ detection of γ , X-rays, e^-



- Interaction only on atomic electrons


→ detection of e.m. radiation



- Interaction of light DMp (LDM) on e^- or nucleus with production of a lighter particle

→ detection of electron/nucleus recoil energy

e.g. sterile ν




... also other ideas ...

e.g. signals from these candidates are **completely lost** in experiments based on "rejection procedures" of the e.m. component of their rate

- ... and more

Investigating the presence of a DM particle component in the galactic halo by the model independent annual modulation signature

Drukier, Freese, Spergel PRD86
Freese et al. PRD88



- $v_{\text{sun}} \sim 232 \text{ km/s}$ (Sun velocity in the halo)
- $v_{\text{orb}} = 30 \text{ km/s}$ (Earth velocity around the Sun)
- $\gamma = \pi/3$
- $\omega = 2\pi/T$ $T = 1 \text{ year}$
- $t_0 = 2^{\text{nd}} \text{ June}$ (when v_{\oplus} is maximum)

$$v_{\oplus}(t) = v_{\text{sun}} + v_{\text{orb}} \cos\gamma \cos[\omega(t-t_0)]$$

$$S_k[\eta(t)] = \int_{\Delta E_k} \frac{dR}{dE_R} dE_R \cong S_{0,k} + S_{m,k} \cos[\omega(t-t_0)]$$

Expected rate in given energy bin changes because of the Earth's motion around the Sun moving in the Galaxy. Therefore it has different peculiarities (e.g. the phase) with respect to those effects connected with seasons

Requirements:

- 1) Modulated rate according cosine
- 2) In a definite low energy range
- 3) With a proper period (1 year)
- 4) With proper phase (about 2nd June)
- 5) For single hit in a multi-detector set-up
- 6) With modulated amplitude in the region of maximal sensitivity < 7% (for usually adopted halo distributions, but it can be larger in case of some possible scenarios)

To mimic this signature, spurious effects and side reactions must not only be able to account for the whole observed modulation amplitude, but also to satisfy contemporaneously all the requirements

DAMA/NaI: ≈ 100 kg NaI(Tl)

Performances: N.Cim.A112(1999)545-575, EPJC18(2000)283,
Riv.N.Cim.26 n. 1(2003)1-73, IJMPD13(2004)2127


Results on rare processes:

- Possible Pauli exclusion principle violation PLB408(1997)439
- CNC processes PRC60(1999)065501
- Electron stability and non-paulian transitions in Iodine atoms (by L-shell) PLB460(1999)235
- Search for solar axions PLB515(2001)6
- Exotic Matter search EPJdirect C14(2002)1
- Search for superdense nuclear matter EPJA23(2005)7
- Search for heavy clusters decays EPJA24(2005)51

Results on DM particles:

- PSD PLB389(1996)757
- Investigation on diurnal effect N.Cim.A112(1999)1541
- Exotic Dark Matter search PRL83(1999)4918
- Annual Modulation Signature

PLB424(1998)195, PLB450(1999)448, PRD61(1999)023512, PLB480(2000)23, EPJC18(2000)283,
PLB509(2001)197, EPJC23(2002)61, PRD66(2002)043503, Riv.N.Cim.26 n.1 (2003)1, IJMPD13(2004)
2127, IJMPA21(2006)1445, EPJC47(2006)263, IJMPA22(2007)3155, EPJC53(2008)205, PRD77(2008)
023506, MPLA23(2008)2125.



model independent evidence of a particle DM component in the galactic halo at 6.3σ C.L.

total exposure (7 annual cycles) 0.29 ton \times yr

Installing the DAMA/LIBRA set-up ~250 kg ULB NaI(Tl)



Residual contaminations in the new DAMA/LIBRA NaI(Tl) detectors: ^{232}Th , ^{238}U and ^{40}K at level of 10^{-12} g/g


- *Radiopurity, performances, procedures, etc.*: NIMA592(2008)297
- *Results on DM particles: Annual Modulation Signature*: EPJC56(2008)333, EPJC67(2010)39
- *Results on rare processes: PEP violation in Na and I*: EPJC62(2009)327

...calibration procedures




DAMA/LIBRA calibrations

Low energy: various external gamma sources (^{241}Am , ^{133}Ba) and internal X-rays or gamma's (^{40}K , ^{125}I , ^{129}I), routine calibrations with ^{241}Am




$$\frac{\sigma_{LE}}{E} = \frac{(0.448 \pm 0.035)}{\sqrt{E(\text{keV})}} + (9.1 \pm 5.1) \cdot 10^{-3}$$

High energy: external sources of gamma rays (e.g. ^{137}Cs , ^{60}Co and ^{133}Ba) and gamma rays of 1461 keV due to ^{40}K decays in an adjacent detector, tagged by the 3.2 keV X-rays




$$\frac{\sigma_{HE}}{E} = \frac{(1.12 \pm 0.06)}{\sqrt{E(\text{keV})}} + (17 \pm 23) \cdot 10^{-4}$$

Thus, here and hereafter keV means keV electron equivalent




The curves superimposed to the experimental data have been obtained by simulations




Noise rejection near the energy threshold

Typical pulse profiles of PMT noise and of scintillation event with the same area, just above the energy threshold of 2 keV

The different time characteristics of PMT noise (decay time of order of tens of ns) and of scintillation event (decay time about 240 ns) can be investigated building several variables




2-4 keV



PMT noise

Scintillation pulses

4-6 keV



From the Waveform Analyser
2048 ns time window:

$$X_1 = \frac{\text{Area (from 100 ns to 600 ns)}}{\text{Area (from 0 ns to 600 ns)}}$$

$$X_2 = \frac{\text{Area (from 0 ns to 50 ns)}}{\text{Area (from 0 ns to 600 ns)}}$$

- The separation between noise and scintillation pulses is very good.
- Very clean samples of scintillation events selected by stringent acceptance windows.
- The related efficiencies evaluated by calibrations with ^{241}Am sources of suitable activity in the same experimental conditions and energy range as the production data (efficiency measurements performed each ~10 days; typically 10^4 - 10^5 events per keV collected)

This is the only procedure applied to the analysed data

Infos about DAMA/LIBRA data taking

Period		Mass (kg)	Exposure (kg × day)	α - β^2
DAMA/LIBRA-1	Sep. 9, 2003 – July 21, 2004	232.8	51405	0.562
DAMA/LIBRA-2	July 21, 2004 – Oct. 28, 2005	232.8	52597	0.467
DAMA/LIBRA-3	Oct. 28, 2005 – July 18, 2006	232.8	39445	0.591
DAMA/LIBRA-4	July 19, 2006 – July 17, 2007	232.8	49377	0.541
DAMA/LIBRA-5	July 17, 2007 – Aug. 29, 2008	232.8	66105	0.468
DAMA/LIBRA-6	Nov. 12, 2008 – Sep. 1, 2009	242.5	58768	0.519
DAMA/LIBRA-1 to -6	Sep. 9, 2003 – Sep. 1, 2009		317697 = 0.87 ton×yr	0.519

- **calibrations: ≈ 72 M events from sources**
- **acceptance window eff: 82 M events (≈ 3 M events/keV)**
- EPJC56(2008)333
- EPJC67(2010)39

DAMA/NaI (7 years) + DAMA/LIBRA (6 years)

total exposure: 425428 kg×day = 1.17 ton×yr

• First upgrade on Sept 2008:

- replacement of some PMTs in HP N₂ atmosphere
- restore 1 detector to operation
- new Digitizers installed (U1063A Acqiris 1GS/s 8-bit High-Speed cPCI)
- new DAQ system with optical read-out installed



Second upgrade on Nov/Dec 2010



all PMTs replaced with new ones of higher Q.E.





Since Dec 2010 data taking and optimizations in this new configuration started

Dark Matter investigation by model-independent annual modulation signature

DAMA/NaI (7 years) + DAMA/LIBRA (6 years). Total exposure: **1.17 ton×yr** EPJC 56(2008)333, EPJC 67(2010)39
 (the **largest** exposure ever collected in this field)

Experimental single-hit residuals rate vs time in 2-6 keV



$A \cos[\omega(t-t_0)]$

continuous lines: $t_0 = 152.5$ d, $T = 1.00$ y

$A = (0.0114 \pm 0.0013)$ cpd/kg/keV

$\chi^2/\text{dof} = 64.7/79$ 8.8σ C.L.

Absence of modulation? No

$\chi^2/\text{dof} = 140/80$ $P(A=0) = 4.3 \times 10^{-5}$


fit with all the parameters free:

$A = (0.0116 \pm 0.0013)$ cpd/kg/keV

$t_0 = (146 \pm 7)$ d

$T = (0.999 \pm 0.002)$ y 8.9σ C.L.

Comparison between **single hit residual rate (red points)** and **multiple hit residual rate (green points)** for (DAMA/LIBRA 1-6); Clear modulation in the single hit events $A = (0.0091 \pm 0.0014)$ cpd/kg/keV; No modulation in the residual rate of the multiple hit events $A = -(0.0006 \pm 0.0004)$ cpd/kg/keV



Multiple hits events = Dark Matter particle "switched off"

No systematics or side reaction able to account for the measured modulation amplitude and to satisfy all the peculiarities of the signature

This result offers an additional strong support for the presence of DM particles in the galactic halo further excluding any side effect either from hardware or from software procedures or from background

The data favor the presence of a modulated behaviour with all the proper features for DM particles in the galactic halo at about 9σ C.L.

Modulation amplitudes measured in each one of the 13 one-year experiments (DAMA/NaI and DAMA/LIBRA)


	A (cpd/kg/keV)	T= 2 π / ω (yr)	t ₀ (day)	C.L.
DAMA/NaI (7 years)				
(2÷4) keV	0.0252 ± 0.0050	1.01 ± 0.02	125 ± 30	5.0 σ
(2÷5) keV	0.0215 ± 0.0039	1.01 ± 0.02	140 ± 30	5.5 σ
(2÷6) keV	0.0200 ± 0.0032	1.00 ± 0.01	140 ± 22	6.3 σ
DAMA/LIBRA (6 years)				
(2÷4) keV	0.0180 ± 0.0025	0.996 ± 0.002	135 ± 8	7.2 σ
(2÷5) keV	0.0134 ± 0.0018	0.997 ± 0.002	140 ± 8	7.4 σ
(2÷6) keV	0.0098 ± 0.0015	0.999 ± 0.002	146 ± 9	6.5 σ
DAMA/NaI + DAMA/LIBRA				
(2÷4) keV	0.0194 ± 0.0022	0.996 ± 0.002	136 ± 7	8.8 σ
(2÷5) keV	0.0149 ± 0.0016	0.997 ± 0.002	142 ± 7	9.3 σ
(2÷6) keV	0.0116 ± 0.0013	0.999 ± 0.002	146 ± 7	8.9 σ

DAMA/NaI (7 annual cycles: 0.29 ton x yr) +
DAMA/LIBRA (6 annual cycles: 0.87 ton x yr)
total exposure: 425428 kg×day = 1.17 ton×yr

A, T, t₀ obtained by fitting the single-hit data with $A\cos[\omega(t-t_0)]$

- The modulation amplitudes for the (2 – 6) keV energy interval, obtained when fixing the period at 1 yr and the phase at 152.5 days, are: (0.019±0.003) cpd/kg/keV for DAMA/NaI and (0.010±0.002) cpd/kg/keV for DAMA/LIBRA.
- Thus, their difference: (0.009±0.004) cpd/kg/keV is $\sim 2\sigma$ which corresponds to a modest, but non negligible probability.

The χ^2 test ($\chi^2 = 9.3, 12.2$ and 10.1 over 12 d.o.f. for the three energy intervals, respectively) and the *run test* (lower tail probabilities of 57%, 47% and 35% for the three energy intervals, respectively) accept at 90% C.L. the hypothesis that the modulation amplitudes are normally fluctuating around their best fit values.



Compatibility among the annual cycles


Energy distribution of the modulation amplitudes

$$R(t) = S_0 + S_m \cos[\omega(t - t_0)]$$

here $T = 2\pi/\omega = 1$ yr and $t_0 = 152.5$ day

DAMA/NaI (7 years) + DAMA/LIBRA (6 years)

total exposure: 425428 kg×day ≈ 1.17 ton×yr




A clear modulation is present in the (2-6) keV energy interval, while S_m values compatible with zero are present just above

The S_m values in the (6-20) keV energy interval have random fluctuations around zero with χ^2 equal to 27.5 for 28 degrees of freedom


Rate behaviour above 6 keV

• No Modulation above 6 keV



Mod. Ampl. (6-10 keV): cpd/kg/keV
 (0.0016 ± 0.0031) DAMA/LIBRA-1
 $-(0.0010 \pm 0.0034)$ DAMA/LIBRA-2
 $-(0.0001 \pm 0.0031)$ DAMA/LIBRA-3
 $-(0.0006 \pm 0.0029)$ DAMA/LIBRA-4
 $-(0.0021 \pm 0.0026)$ DAMA/LIBRA-5
 (0.0029 ± 0.0025) DAMA/LIBRA-6
 → statistically consistent with zero

DAMALIBRA-1 to -6



$\sigma \approx 1\%$, fully accounted by statistical considerations

• No modulation in the whole energy spectrum: studying integral rate at higher energy, R_{90}

- R_{90} percentage variations with respect to their mean values for single crystal in the DAMA/LIBRA running periods
- Fitting the behaviour with time, adding a term modulated with period and phase as expected for DM particles:

consistent with zero

Period	Mod. Ampl.
DAMA/LIBRA-1	$-(0.05 \pm 0.19)$ cpd/kg
DAMA/LIBRA-2	$-(0.12 \pm 0.19)$ cpd/kg
DAMA/LIBRA-3	$-(0.13 \pm 0.18)$ cpd/kg
DAMA/LIBRA-4	(0.15 ± 0.17) cpd/kg
DAMA/LIBRA-5	(0.20 ± 0.18) cpd/kg
DAMA/LIBRA-6	$-(0.20 \pm 0.16)$ cpd/kg

+ if a modulation present in the whole energy spectrum at the level found in the lowest energy region → $R_{90} \sim \text{tens cpd/kg} \rightarrow \sim 100 \sigma$ far away

No modulation above 6 keV
 This accounts for all sources of bckg and is consistent with studies on the various components

The analysis at energies above 6 keV, the analysis of the multiple-hits events and the statistical considerations about S_m already exclude any sizable presence of systematical effects

Additional investigations on the stability parameters

Modulation amplitudes obtained by fitting the time behaviours of main running parameters, acquired with the production data, when including a DM-like modulation

Running conditions stable at a level better than 1% also in the two new running periods

	DAMA/LIBRA-1	DAMA/LIBRA-2	DAMA/LIBRA-3	DAMA/LIBRA-4	DAMA/LIBRA-5	DAMA/LIBRA-6
Temperature	$(-0.0001 \pm 0.0061) \text{ }^\circ\text{C}$	$(0.0026 \pm 0.0086) \text{ }^\circ\text{C}$	$(0.001 \pm 0.015) \text{ }^\circ\text{C}$	$(0.0004 \pm 0.0047) \text{ }^\circ\text{C}$	$(0.0001 \pm 0.0036) \text{ }^\circ\text{C}$	$(0.0007 \pm 0.0059) \text{ }^\circ\text{C}$
Flux N_2	$(0.13 \pm 0.22) \text{ l/h}$	$(0.10 \pm 0.25) \text{ l/h}$	$(-0.07 \pm 0.18) \text{ l/h}$	$(-0.05 \pm 0.24) \text{ l/h}$	$(-0.01 \pm 0.21) \text{ l/h}$	$(-0.01 \pm 0.15) \text{ l/h}$
Pressure	$(0.015 \pm 0.030) \text{ mbar}$	$(-0.013 \pm 0.025) \text{ mbar}$	$(0.022 \pm 0.027) \text{ mbar}$	$(0.0018 \pm 0.0074) \text{ mbar}$	$(-0.08 \pm 0.12) \times 10^{-2} \text{ mbar}$	$(0.07 \pm 0.13) \times 10^{-2} \text{ mbar}$
Radon	$(-0.029 \pm 0.029) \text{ Bq/m}^3$	$(-0.030 \pm 0.027) \text{ Bq/m}^3$	$(0.015 \pm 0.029) \text{ Bq/m}^3$	$(-0.052 \pm 0.039) \text{ Bq/m}^3$	$(0.021 \pm 0.037) \text{ Bq/m}^3$	$(-0.028 \pm 0.036) \text{ Bq/m}^3$
Hardware rate above single photoelectron	$(-0.20 \pm 0.18) \times 10^{-2} \text{ Hz}$	$(0.09 \pm 0.17) \times 10^{-2} \text{ Hz}$	$(-0.03 \pm 0.20) \times 10^{-2} \text{ Hz}$	$(0.15 \pm 0.15) \times 10^{-2} \text{ Hz}$	$(0.03 \pm 0.14) \times 10^{-2} \text{ Hz}$	$(0.08 \pm 0.11) \times 10^{-2} \text{ Hz}$

All the measured amplitudes well compatible with zero

+ none can account for the observed effect


(to mimic such signature, spurious effects and side reactions must not only be able to account for the whole observed modulation amplitude, but also simultaneously satisfy all the 6 requirements)

Summary of the results obtained in the additional investigations of possible systematics or side reactions

(previous exposure and details see: NIMA592(2008)297, EPJC56(2008)333, arXiv:0912.4200, arXiv:1007.0595)

DAMA/LIBRA 1-6

<i>Source</i>	<i>Main comment</i>	<i>Cautious upper limit (90% C.L.)</i>
RADON	Sealed Cu box in HP Nitrogen atmosphere, 3-level of sealing, etc.	$<2.5 \times 10^{-6}$ cpd/kg/keV
TEMPERATURE	Installation is air conditioned+ detectors in Cu housings directly in contact with multi-ton shield → huge heat capacity + T continuously recorded	$<10^{-4}$ cpd/kg/keV
NOISE	Effective full noise rejection near threshold	$<10^{-4}$ cpd/kg/keV
ENERGY SCALE	Routine + intrinsic calibrations	$<1-2 \times 10^{-4}$ cpd/kg/keV
EFFICIENCIES	Regularly measured by dedicated calibrations	$<10^{-4}$ cpd/kg/keV
BACKGROUND	No modulation above 6 keV; no modulation in the (2-6) keV <i>multiple-hits</i> events; this limit includes all possible sources of background	$<10^{-4}$ cpd/kg/keV
SIDE REACTIONS	Muon flux variation measured at LNGS	$<3 \times 10^{-5}$ cpd/kg/keV



+ they cannot satisfy all the requirements of annual modulation signature




Thus, they cannot mimic the observed annual modulation effect

The μ case

MonteCarlo simulation


- muon intensity distribution
- Gran Sasso rock overburden map

events where just one detector fires



DAMA/LIBRA surface $\approx 0.15 \text{ m}^2$
 μ flux @ DAMA/LIBRA $\approx 2.5 \mu/\text{day}$

1.



Case of fast neutrons produced by μ

Φ_μ @ LNGS $\approx 20 \mu \text{ m}^{-2} \text{ d}^{-1}$ ($\pm 2\%$ modulated)
 Measured neutron Yield @ LNGS: $Y = 1 \div 7 \cdot 10^{-4} \text{ n}/\mu / (\text{g}/\text{cm}^2)$
 $R_n = (\text{fast n by } \mu) / (\text{time unit}) = \Phi_\mu Y M_{\text{eff}}$

Annual modulation amplitude at low energy due to μ modulation:

$$S_m^{(\mu)} = R_n g \varepsilon f_{\Delta E} f_{\text{single}} 2\% / (M_{\text{setup}} \Delta E)$$

g = geometrical factor; ε = detection effc. by elastic scattering
 $f_{\Delta E}$ = energy window ($E > 2 \text{ keV}$) effc.; f_{single} = single hit effc.

Hyp.: $M_{\text{eff}} = 15 \text{ tons}$; $g \approx \varepsilon \approx f_{\Delta E} \approx f_{\text{single}} \approx 0.5$ (cautiously)
 Knowing that: $M_{\text{setup}} \approx 250 \text{ kg}$ and $\Delta E = 4 \text{ keV}$

$$S_m^{(\mu)} < (0.4 \div 3) \times 10^{-5} \text{ cpd}/\text{kg}/\text{keV}$$

Moreover, this modulation also induces a variation in other parts of the energy spectrum and in the *multi-hits* events

It cannot mimic the signature: already excluded also by R_{90} , by *multi-hits* analysis + different phase, etc.

Can (whatever) hypothetical cosmogenic products be considered as side effects, assuming that they might produce:

- only events at low energy,
- only *single-hit* events,
- no sizable effect in the *multiple-hit* counting rate
- pulses with time structure as scintillation light

?

But, its phase should be (much) larger than μ phase, t_μ :

- if $\tau \ll T/2\pi$: $t_{\text{side}} = t_\mu + \tau$
- if $\tau \gg T/2\pi$: $t_{\text{side}} = t_\mu + T/4$

It cannot mimic the signature: different phase

The phase of the muon flux at LNGS is roughly around middle of July and largely variable from year to year. Last meas. by LVD and BOREXINO partially overlapped with DAMA/NaI and fully with DAMA/LIBRA: 1.5% modulation and phase LVD = July 5th \pm 15 d, BOREXINO = July 6th \pm 6 d

DAMA/NaI + DAMA/LIBRA measured a stable phase: May, 26th \pm 7 days

This phase is 7.1 σ far from July 15th and is 5.7 σ far from July 6th

R_{90} , multi-hits, phase, and other analyses

NO

more about the phase of muons ...

2.

The DAMA: modulation amplitude 10^{-2} cpd/kg/keV, in 2-6 keV energy range for single hit events; phase:

May 26 \pm 7 days


(stable over 13 years)

μ flux @ LNGS (MACRO, LVD, BOREXINO)
 $\approx 3 \cdot 10^{-4} \text{ m}^{-2}\text{s}^{-1}$; modulation amplitude 1.5%; phase:

July 6 \pm 6 days (BOREXINO, CSN2 sept. 2010)

but

the muon phase differs from year to year (error no purely statistical); LVD/BOREXINO phase value is a “mean” of the muon phase of each year



The DAMA phase is 5.7σ far from the LVD/BOREXINO phases of muons (7.3σ far from MACRO measured phase)

- 1) if we assume for a while that the real value of the DAMA phase is June 16th (that is 3σ fluctuation from the measured value), it is well far from all the measured phases of muons by LVD, MACRO and BOREXINO, in all the years
- 2) Moreover, considering the seasonal weather condition in Gran Sasso, it is quite impossible that the maximum temperature of the outer atmosphere (on which μ flux modulation is dependent) is observed in the middle of June

Inconsistency of the phase between DAMA signal and μ modulation

Summarizing

- Presence of modulation for 13 annual cycles at 8.9σ C.L. with the proper distinctive features of the DM signature; all the features satisfied by the data over 13 independent experiments of 1 year each one
- The total exposure by former DAMA/NaI and present DAMA/LIBRA is **1.17 ton × yr (13 annual cycles)**
- In fact, as required by the DM annual modulation signature:

1. The *single-hit* events show a clear cosine-like modulation, as expected for the DM signal

2. Measured period is equal to (0.999 ± 0.002) yr, well compatible with the 1 yr period, as expected for the DM signal

3. Measured phase (146 ± 7) days is well compatible with 152.5 days, as expected for the DM signal

4. The modulation is present only in the low energy (2-6) keV interval and not in other higher energy regions, consistently with expectation for the DM signal

5. The modulation is present only in the *single-hit* events, while it is absent in the *multiple-hits*, as expected for the DM signal



6. The measured modulation amplitude in NaI(Tl) of the *single-hit* events in (2-6) keV is: (0.0116 ± 0.0013) cpd/kg/keV (8.9σ C.L.).

No systematic or side process able to simultaneously satisfy all the many peculiarities of the signature and to account for the whole measured modulation amplitude is available



Just few examples of interpretation of the annual modulation in terms of candidate particles in some scenarios

WIMP: SI



- Not best fit
- About the same C.L.



WIMP: SI & SD $\theta = 2.435$



LDM, bosonic DM



EPJC56(2008)333

Compatibility with several candidates; other ones are open

About interpretation

See e.g.: Riv.N.Cim.26 n.1(2003)1,
IJMPD13(2004)2127, EPJC47(2006)263,
IJMPA21(2006)1445, EPJC56(2008)333,
arXiv:1106.4667 in press on PRD

...models...

- Which particle?
- Which interaction coupling?
- Which Form Factors for each target-material?
- Which Spin Factor?
- Which nuclear model framework?
- Which scaling law?
- Which halo model, profile and related parameters?
- Streams?
- ...

...and experimental aspects...

- Exposures
- Energy threshold
- Detector response(phe/keV)
- Energy scale and energy resolution
- Calibrations
- Stability of all the operating conditions.
- Selections of detectors and of data.
- Subtraction/rejection procedures and stability in time of all the selected windows and related quantities
- Efficiencies
- Definition of fiducial volume and non-uniformity
- Quenching factors, channeling
- ...

Uncertainty in experimental parameters, as well as necessary assumptions on various related astrophysical, nuclear and particle-physics aspects, affect all the results at various extent, both in terms of exclusion plots and in terms of allowed regions/volumes. Thus comparisons with a fixed set of assumptions and parameters' values are intrinsically strongly uncertain.


... an example in literature...

Supersymmetric expectations in MSSM

- Assuming for the neutralino a dominant purely SI coupling
- when releasing the gaugino mass unification at GUT scale: $M_1/M_2 \neq 0.5$ ($<$);
(where M_1 and M_2 U(1) and SU(2) gaugino masses)

Relic neutralino in effMSSM


N. Fornengo, S. Scopel, A. Bottino (2010)



DAMA allowed regions for a particular set of astrophysical, nuclear and particle Physics assumptions with and without channeling


CoGeNT and CRESST

If the two CDMS events are interpreted as relic neutralino interactions



DAMA allowed regions for a particular set of astrophysical, nuclear and particle Physics assumptions without (green), with (blue) channeling, with en.dep. Q.F.(red)

CoGeNT



... examples in some given frameworks

DM particle with preferred inelastic interaction

- In the **Inelastic DM (iDM)** scenario, WIMPs scatter into an excited state, split from the ground state by an energy comparable to the available kinetic energy of a Galactic WIMP.



→ W has two mass states χ^+ , χ^- with δ mass splitting

→ Kinematical constraint for iDM

$$\frac{1}{2} \mu v^2 \geq \delta \Leftrightarrow v \geq v_{thr} = \sqrt{\frac{2\delta}{\mu}}$$

DAMA/NaI+DAMA/LIBRA

Slices from the 3-dimensional allowed volume

iDM interaction on Iodine nuclei

Fund. Phys. 40(2010)900


iDM interaction on Tl nuclei of the NaI(Tl) dopant?

- For **large splittings**, the dominant scattering in NaI(Tl) can occur off of **Thallium nuclei**, with $A \sim 205$, which are present as a dopant at the 10^{-3} level in NaI(Tl) crystals.

arXiv:1007.2688

- Inelastic scattering WIMPs with **large splittings** do not give rise to sizeable contribution on Na, I, Ge, Xe, Ca, O, ... nuclei.

... and more considering experimental and theoretical uncertainties



Model-independent evidence by DAMA/NaI and DAMA/LIBRA

well compatible with several candidates


(in many possible astrophysical, nuclear and particle physics scenarios)

Some other papers on compatibility among results: **Next-to-minimal models** (JCAP0908(2009)032, PRD79(2009)023510, JCAP0706(2007)008, arXiv:1009.2555,1009.0549), **Inelastic DM** (PRD79(2009)043513), **Sneutrino DM** (JHEP0711(2007)029, arXiv:1105.4878), **Resonant DM** (arXiv:0909.2900), **Cogent results** (arXiv:1002.4703,1106.0650), **DM from exotic 4th generation quarks** (arXiv:1002.3366), **Light WIMP DM** (PRD81(2010)107302, PRD83(2011)015001, arXiv:1003.0014,1007.1005,1106.1066), **Composite DM** (IJMPD19(2010)1385), **Light scalar WIMP through Higgs portal** (PRD82(2010)043522, JCAP0810(2010)034), **Specific two higgs doublet models** (arXiv:1106.3368), **exothermic DM** (arXiv:1004.0937), **Secluded WIMPs** (PRD79(2009)115019), **iDM on TI** (arXiv:1007.2688), **Mirror DM** (arXiv:1001.0096,1106.2688, PRD82(2010)095001, JCAP1107(2011)009, JCAP1009(2010)022), **Asymmetric DM** (arXiv:1105.5431), **Isospin-violating models** (JCAP1008(2010)018, arXiv:1102.4331,1105.3734), **Singlet DM** (JHEP0905(2009)036, arXiv:1011.6377), **Specific GU** (arXiv:1106.3583), and more (arXiv:1105.5121,1105.3734,1011.1499, JCAP1008(2010)018, PRD82(2010)115019)...

Possible model dependent positive hints from Indirect searches (but interpretation, evidence itself, derived mass and cross sections depend e.g. on bckg modeling, on DM spatial velocity distribution in the galactic halo, etc.)

not in conflict with DAMA results, null results not in conflict as well

Available results from direct searches using different target materials and approaches **do not give any robust conflict & compatibility with positive excesses**



WHAT NEXT?

- Upgrade in fall 2010 concluded: all PMTs replaced with new ones of higher Q.E. to lower the software energy threshold and improve general features.
- New preamplifiers and trigger modules realised to further implement low energy studies.
- Suitable exposure planned in the new running conditions to deeper study the nature of the particles and features of related astrophysical, nuclear and particle physics aspects.
- A possible 1 ton ULB NaI(Tl) set-up experiment proposed by DAMA in 1996

Conclusions

- **Positive evidence for the presence of DM particles in the galactic halo now supported at 8.9σ C.L. (cumulative exposure $1.17 \text{ ton}\times\text{yr}$ – 13 annual cycles DAMA/NaI and DAMA/LIBRA)**
- **Updating of corollary analyses in some of the many possible scenarios for DM candidates, interactions, halo models, nuclear/atomic properties, etc. is in progress.**
- **Upgrade in fall 2010 concluded and further improvements foreseen.**
- **Analyses/data taking to investigate other rare processes in progress**

DAMA/LIBRA still the highest radiopure set-up in the field with the largest sensitive mass, full control of running conditions, the largest duty-cycle, exposure orders of magnitude larger than any other activity in the field, etc.